

Homecare-Root Canal Therapy

Now that you've had root canal therapy, it's important to follow these recommendations to ensure healing.

Caring for a temporary restoration

If we placed a temporary filling or crown on your tooth, avoid chewing for at least one-half hour to allow the restoration to harden.

To keep your temporary restoration in place, avoid eating hard or sticky food, especially chewing gum. If possible, chew only on the opposite side of your mouth.

It's not a problem for a small portion of a temporary filling to wear away or break off, but if the entire filling wears out, or if a temporary crown comes off, call us so that it can be replaced.

Don't eat if your mouth is numb.

Take medications as directed.

Continue to brush and floss normally.

Chewing and eating

If we used an anesthetic during the procedure, avoid chewing until the numbness has worn off completely. Your lips, teeth, and tongue may be numb for several hours.

Brushing and flossing

Brush and floss normally.

Medications and discomfort

If antibiotics were prescribed, continue to take them for the indicated length of time, even if all symptoms and signs of infection are gone.

To control discomfort, take pain medication before the anesthetic has worn off or as recommended. It's normal to experience some discomfort for several days after a root canal appointment, especially when chewing.

To further reduce discomfort or swelling, rinse your mouth three times a day with warm salt water. Use about one teaspoon of salt per glass of warm water.

When to call us

Call our office if your bite feels uneven, you have sensitivity or discomfort that increases or continues beyond 3 or 4 days, your temporary filling or crown comes off, or you have any questions or concerns.